

University of Groningen

The genes for secretion and maturation of lactococcins are located on the chromosome of *Lactococcus lactis* IL1403

Venema, Koen; Dost, Michiel H.R.; Beun, Philip A.H.; Haandrikman, Alfred J.; Venema, Gerard; Kok, Jan

Published in:
Applied and environmental microbiology

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
1996

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Venema, K., Dost, M. H. R., Beun, P. A. H., Haandrikman, A. J., Venema, G., & Kok, J. (1996). The genes for secretion and maturation of lactococcins are located on the chromosome of *Lactococcus lactis* IL1403. *Applied and environmental microbiology*, 62(10), 1689-1692.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

The Genes for Secretion and Maturation of Lactococcins Are Located on the Chromosome of *Lactococcus lactis* IL1403

KOEN VENEMA,[†] MICHIEL H. R. DOST, PHILIP A. H. BEUN, ALFRED J. HAANDRIKMAN,[‡]
GERARD VENEMA,* AND JAN KOK

Department of Genetics, Groningen Biomolecular Sciences and Biotechnology Institute,
University of Groningen, 9751 NN Haren, The Netherlands

Received 27 July 1995/Accepted 11 February 1996

Southern hybridization and PCR analysis were used to show that *Lactococcus lactis* IL1403, a plasmid-free strain that does not produce bacteriocin, contains genes on its chromosome that are highly homologous to *lcnC* and *lcnD* and encode the lactococcin secretion and maturation system. The *lcnC* and *lcnD* homologs on the chromosome of IL1403 were interrupted independently by Campbell-type integrations. Both insertion mutants were unable to secrete active lactococcin. Part of the chromosomal *lcnC* gene was cloned and sequenced. Only a few nucleotide substitutions occurred, compared with the plasmid-encoded *lcnC* gene, and these did not lead to changes in the deduced amino acid sequence. No genes homologous to those for lactococcin A, B, or M could be detected in IL1403, and the strain does not produce bacteriocin activity.

Lactococcins are bacteriocins produced by *Lactococcus lactis*. van Belkum et al. (29, 30) determined the nucleotide sequences of three bacteriocin determinants from the lactococcal plasmid p9B4-6. In detailed mutational analyses, the lactococcin structural and immunity genes were identified (29, 30). Conclusive evidence that *lcnA* is indeed the structural gene of lactococcin A came from Holo et al. (10). By comparing the amino acid sequence determined from the purified bacteriocin with that deduced from the structural gene, it was evident that lactococcin A was synthesized as a 75-amino-acid precursor with an N-terminal extension of 21 amino acids. Processing of this precursor takes place behind a glycine doublet. Recently, LcnB has also been purified and its N-terminal amino acid sequence has been determined (31). This bacteriocin is also processed after a double glycine. The glycine processing site is present in many class II peptide bacteriocins of lactic acid bacteria (13), even in some lantibiotics, and in colicin V, an *Escherichia coli* peptide bacteriocin (5, 7, 8, 11, 17, 19, 27).

lcnA is also present on pNP2 of *L. lactis* subsp. *lactis* bv. diacetylactis WM4 (25), and by Tn5 transposon mutagenesis, it was shown that two additional genes are required for bacteriocin activity. These genes, designated *lcnC* and *lcnD*, are located in an operon immediately upstream of the lactococcin A structural and immunity genes (25). Comparisons of LcnC and LcnD with protein database sequences revealed that these proteins have significant similarities to gram-negative proteins implicated in signal sequence-independent secretion pathways (2, 9, 24). It is now generally accepted that LcnC and LcnD are required for the secretion of lactococcins via a system dedicated to bacteriocin export (14, 25). Similarly dedicated secretion systems have now been proposed for other lactic acid bacterium bacteriocins (15, 17).

In retrospect, it is surprising that lactococcin production and secretion were found with plasmids which carry only the lactococcin structural and immunity genes but do not encode the postulated secretion function. Holo et al. (10) were able to

detect only reduced production of lactococcin A in *L. lactis* IL1403. This strain was also used throughout the works of van Belkum et al. (28–30). In all other strains, no lactococcin production was detectable. In this study, we investigated why strain IL1403 is capable of producing and externalizing active lactococcins in the absence of the plasmid-encoded *lcnC* and *lcnD* genes.

MATERIALS AND METHODS

Bacterial strains, plasmids, and growth conditions. The bacterial strains and plasmids used in this study are listed in Table 1. *L. lactis* was cultured in M17 broth (26) supplemented with 0.5% glucose. Agar (1.5%) was added to plates. *E. coli* was cultured on TY medium (21). Erythromycin and chloramphenicol were used at final concentrations of 5 µg/ml each for *L. lactis* and of 100 and 10 µg/ml, respectively, for *E. coli*.

Molecular cloning and DNA sequencing. Transformations of *L. lactis* and *E. coli* were done as described earlier (32). General DNA cloning and manipulation techniques and Southern hybridizations were carried out essentially as described by Sambrook et al. (22). Probes for Southern hybridizations were labelled with an ECL labelling kit (Amersham, Little Chalfont, Buckinghamshire, United Kingdom). DNA sequencing was done on double-stranded DNA by the dideoxy chain termination method (23), using a T7 sequencing kit and the manufacturer's (Pharmacia, Uppsala, Sweden) protocol.

Bacteriocin assay. The bacteriocin spot test was used to determine bacteriocin activity, as described before (31). One arbitrary unit was defined as the reciprocal of the highest dilution forming a visible halo. Overlay assays on colonies were done as described before (31). *L. lactis* IL1403 was used as the lactococcin indicator strain throughout this study.

Construction of plasmids. Internal fragments from the plasmid-derived *lcnC* and *lcnD* genes (*Hind*III-*Nsi*I and *Hinc*II-*Hae*III fragments for *lcnC* and *lcnD*, respectively) were cloned in pORI28 (16) to give pINT4C and pINT1D, respectively. These plasmids were maintained in the *E. coli* helper strain EC1000, which contains on its chromosome a copy of *repA*, encoding the pWV01 replication protein RepA. The plasmids from this strain were isolated and used to transform *L. lactis* IL1403. Campbell-type integrants were selected on plates containing erythromycin, as described by Leenhouts and Venema (16).

pHVB was constructed to test bacteriocin production in integrants. This plasmid is a theta-replicating plasmid derived from pAMβ1 and does not interfere with the integrated state of the pINT plasmids. To obtain pHVB, pMB580 was cut with *Eco*RI and *Xba*I. The fragment carrying *lcnB* was isolated and cloned into pHV1432 cut with the same restriction endonucleases.

PCR. PCR was performed according to standard protocols (22) with *Taq* polymerase (Boehringer Mannheim GmbH, Mannheim, Germany). The primers used for PCR were synthesized with model 381A DNA synthesizer (Applied Biosystems, Foster City, Calif.).

Plasmid rescue. Chromosomal DNA of *L. lactis* IL1403::pINT4C was isolated and cut with various restriction endonucleases. After self-ligation, the DNA was used to transform *E. coli* EC1000, with selection for erythromycin resistance.

* Corresponding author.

[†] Present address: Dept. of Food Science, North Carolina State University, Box 7624, Raleigh, NC 27695.

[‡] Present address: Hercules European Research Centre, 3771 ME Barneveld, The Netherlands.

TABLE 1. Bacterial strains and plasmids used in this study

Bacterial strain or plasmid	Relevant properties ^a	Source or reference
Bacteria		
<i>L. lactis</i>		
IL1403	Plasmid-free strain; indicator strain for lactococcin B	1
MG1363	Plasmid-free strain	6
<i>E. coli</i> EC1000	Helper strain; MC1000 containing pWV01 <i>repA</i> on chromosome	Laboratory collection
Plasmids		
pMB580	Em ^r ; containing the lactococcin B operon	30
pMB551	Em ^r ; containing most of <i>lcnC</i> and <i>lcnD</i> ; used as probe in Southern hybridization	30
pHV1432	Cm ^r ; cloning vector; pAM β 1 derivative	21
pORI28	Em ^r ; cloning vector lacking <i>repA</i>	16
pINT4C	Em ^r ; integration vector; pORI28 with an internal fragment of <i>lcnC</i>	This work
pINT1D	Em ^r ; integration vector; pORI28 with an internal fragment of <i>lcnD</i>	This work
pHVB	Cm ^r ; pHV1432 derivative containing lactococcin B operon	This work
pKV4	Em ^r ; pIL253 derivative containing <i>lcnCDA</i> and <i>lciA</i>	30a

^a Em^r, erythromycin resistance; Cm^r, chloramphenicol resistance.

RESULTS AND DISCUSSION

The fact that *lcnC* and *lcnD* are essential for extracellular lactococcin activity (25), combined with the observation that in *L. lactis* IL1403 these genes need not be present on a plasmid together with the *lcnA* structural gene (10, 28, 30), prompted us to investigate whether IL1403, a plasmid-free strain, carries genes homologous to *lcnC* and *lcnD* on its chromosome. A DNA fragment encompassing most of plasmid-derived *lcnC* and *lcnD* was used as the probe in Southern hybridizations. A signal was indeed obtained with the chromosome of IL1403 (Fig. 1B, lanes 6 to 9). Subfragments of *lcnC* and *lcnD* also reacted in Southern hybridizations (data not shown). In PCRs on IL1403 chromosomal DNA with primers derived from and spanning almost the entire plasmid-encoded *lcnC* and *lcnD* region (3, 4), fragments with sizes similar to those expected from plasmid-borne *lcnC* and *lcnD* were obtained (Fig. 2A). No PCR products were obtained with primers specific for the 3' end of *lcnD*. By using DNA probes encompassing this part of the gene in a Southern hybridization, a signal was found,

however, suggesting that a homologous region is preserved on the chromosome of IL1403. Apparently, the nucleotide sequence of the part of *lcnD* against which the primers were directed differs significantly from that of the plasmid-encoded gene, preventing primer annealing. Taken together, these results indicate that the chromosome of strain IL1403 contains




FIG. 1. (A) Agarose (0.8%) gel; (B) corresponding Southern hybridization film. Lanes 1, plasmid pMB551 (positive control); lanes 2 to 5, chromosomal DNA of *L. lactis* MG1363 cut with *EcoRV*, *HindIII*, *HpaII*, and *PvuII*, respectively; lanes 6 to 9, chromosomal DNA of *L. lactis* IL1403 cut with *EcoRV*, *HindIII*, *HpaII*, and *PvuII*, respectively; lanes 10, SPP1 marker DNA cut with *EcoRI*. Plasmid pMB551 was used as the probe.




FIG. 2. (A) Schematic representation of the PCR products obtained from the chromosome of IL1403. Schematic outlines for the Campbell-type integration of pINT4C into the chromosome of IL1403 to produce an insertion mutant of *lcnC* (B) and plasmid rescue (C). ORI⁺, origin of replication; Em^r, erythromycin resistance marker; X and Y, restriction endonuclease sites; P, promoter; open box, internal fragment of *lcnC*; hatched box, chromosomally located gene.

9. Higgins, C. F. 1993. ABC transporters: from microorganisms to man. *Annu. Rev. Cell Biol.* **8**:67–113.
10. Holo, H., O. Nilssen, and I. F. Nes. 1991. Lactococcin A, a new bacteriocin from *Lactococcus lactis* subsp. *cremoris*: isolation and characterization of the protein and its gene. *J. Bacteriol.* **173**:3879–3887.
11. Hynes, W. L., J. J. Ferreti, and J. R. Tagg. 1993. Cloning the gene encoding streptococcin A-FF22, a novel lantibiotic produced by *Streptococcus pyogenes*, and determination of its nucleotide sequence. *Appl. Environ. Microbiol.* **59**:1969–1971.
12. Jannière, L., C. Bruand, and S. D. Ehrlich. 1990. Structurally stable *Bacillus subtilis* cloning vectors. *Gene* **87**:53–61.
13. Klaenhammer, T. R. 1993. Genetics of bacteriocins produced by lactic acid bacteria. *FEMS Microbiol. Rev.* **12**:39–86.
14. Kok, J., H. Holo, M. J. van Belkum, A. J. Haandrikman, and I. F. Nes. 1993. Nonnisin bacteriocins in lactococci: biochemistry, genetics and mode of action, p. 121–150. *In* D. Hoover and L. Steenson (ed.), *Bacteriocins of lactic acid bacteria*. Academic Press, New York.
15. Kuipers, O. P., M. M. Beerthuyzen, R. J. Siezen, and W. M. de Vos. 1993. Characterization of the nisin gene cluster *nisABTCIPR* of *Lactococcus lactis*: requirement of expression of *nisA* and *nisI* genes for producer immunity. *Eur. J. Biochem.* **216**:281–291.
16. Leenhouts, K. J., and G. Venema. 1993. Lactococcal plasmid vectors, p. 66–94. *In* K. G. Hardy (ed.), *Plasmids. A practical approach*. Oxford University Press, New York.
17. Marugg, J. D., C. F. Gonzalez, B. S. Kunka, A. M. Ledebøer, M. J. Pucci, M. Y. Toonen, S. A. Walker, L. C. M. Zoetmulder, and P. A. Vandenberg. 1992. Cloning, expression, and nucleotide sequence of genes involved in production of pediocin PA-1, a bacteriocin from *Pediococcus acidilactici* PAC1.0. *Appl. Environ. Microbiol.* **58**:2360–2367.
18. Piard, J.-C., O. P. Kuipers, H. S. Rollet, M. J. Desmazeaud, and W. M. de Vos. 1993. Structure, organization, and expression of the *lct* gene for lactacin 481, a novel lantibiotic produced by *Lactococcus lactis*. *J. Biol. Chem.* **268**:16361–16368.
19. Quadri, L. E. N., M. Sailer, K. L. Roy, J. C. Vederas, and M. E. Stiles. 1994. Chemical and genetic characterization of bacteriocins produced by *Carnobacterium piscicola* LV17B. *J. Biol. Chem.* **269**:12204–12211.
20. Rince, A., A. Dufour, S. Le Pogam, D. Thuault, C. M. Bourgeois, and J. P. Le Pennec. 1994. Cloning, expression, and nucleotide sequence of genes involved in production of lactococcin DR, a bacteriocin from *Lactococcus lactis* subsp. *lactis*. *Appl. Environ. Microbiol.* **60**:1652–1657.
21. Rottlander, E., and T. A. Trautner. 1970. Genetic and transfection studies with *Bacillus subtilis* phage SP50. *Mol. Gen. Genet.* **108**:47–60.
22. Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. *Molecular cloning: a laboratory manual*, 2nd ed. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
23. Sanger, F., S. Nicklen, and A. R. Coulson. 1977. DNA sequencing with chain-terminating inhibitors. *Proc. Natl. Acad. Sci. USA* **74**:5463–5467.
24. Schüle, R., I. Gentschev, H.-J. Mollenkopf, and W. Goebel. 1992. A topological model for the haemolysin translocator protein HlyD. *Mol. Gen. Genet.* **234**:155–163.
25. Stoddard, G. W., J. P. Petzel, M. J. van Belkum, J. Kok, and L. L. McKay. 1992. Molecular analyses of the lactococcin A gene cluster from *Lactococcus lactis* subsp. *lactis* biovar diacetylactis WM4. *Appl. Environ. Microbiol.* **58**:1952–1961.
26. Terzaghi, B. E., and W. E. Sandine. 1975. Improved medium for lactic streptococci and their bacteriophages. *Appl. Microbiol.* **29**:807–813.
27. Tichaczek, P. S., J. Nissen-Meyer, I. F. Nes, R. F. Vogel, and W. P. Hammes. 1992. Characterization of the bacteriocins curvacin A from *Lactobacillus curvatus* LTH1174 and sakacin P from *Lactobacillus sake* LTH673. *Syst. Appl. Microbiol.* **15**:460–468.
28. van Belkum, M. J., B. J. Hayema, A. Geis, J. Kok, and G. Venema. 1989. Cloning of two bacteriocin genes from a lactococcal bacteriocin plasmid. *Appl. Environ. Microbiol.* **55**:1187–1191.
29. van Belkum, M. J., B. J. Hayema, R. E. Jeeninga, J. Kok, and G. Venema. 1991. Organization and nucleotide sequences of two lactococcal bacteriocin operons. *Appl. Environ. Microbiol.* **57**:492–498.
30. van Belkum, M. J., J. Kok, and G. Venema. 1992. Cloning, sequencing, and expression in *Escherichia coli* of *lcnB*, a third bacteriocin determinant from the lactococcal bacteriocin plasmid p9B4-6. *Appl. Environ. Microbiol.* **58**:572–577.
- 30a. Venema, K. Ph.D. thesis, University of Groningen, Groningen, The Netherlands.
31. Venema, K., M. L. Chikindas, J. F. M. L. Seegers, A. J. Haandrikman, K. J. Leenhouts, G. Venema, and J. Kok. General, high yield, and rapid purification method for small, hydrophobic, cationic bacteriocins. Purification of lactococcin B from *Lactococcus lactis* subsp. *cremoris* 9B4 and pediocin PA-1 from *Pediococcus acidilactici* PAC1.0. Submitted for publication.
32. Venema, K., R. E. Haverkort, T. Abec, A. J. Haandrikman, K. J. Leenhouts, L. de Leij, G. Venema, and J. Kok. 1994. Mode of action of LciA, the lactococcin A immunity protein. *Mol. Microbiol.* **14**:521–532.
33. Venema, K., J. Kok, J. D. Marugg, M. Y. Toonen, A. M. Ledebøer, G. Venema, and M. L. Chikindas. 1995. Functional analysis of the pediocin operon of *Pediococcus acidilactici* PAC1.0: PedB is the immunity protein and PedD is the processing enzyme. *Mol. Microbiol.* **17**:515–522.